ARKANSAS
STATE PLANT
BOARD

Darryl Little
Director

P.O. Box 1069 ■ Little Rock, Arkansas 72203
www.plantboard.org
Phone (501) 225-1598
Fax (501) 219-1697

June 11, 2012

To whom it may concern:

Please find enclosed a Section 18 Notice of Specific Exemption along with forms with which to Report Section 18 Applications.

The notice is for the use of Transform WG for Control of Tarnished Plant Bug (Lygus lineolaris) in Cotton.

If you have any questions feel free to contact me (501) 225-1598.

Sincerely,

Brandi Reynolds

Brandi Reynolds
Assistant Director
Pesticide Division

ARKANSAS
STATE PLANT
BOARD

P.O. Box 1069 ■ Little Rock, Arkansas 72203
www.plantboard.org
Phone (501) 225-1598
Fax (501) 219-1697

Darryl Little
Director

NOTICE OF SPECIFIC EXEMPTION FOR THE USE OF TRANSFORM WG FOR CONTROL OF TARNISHED PLANT BUG IN COTTON

June 11, 2012

The Environmental Protection Agency has granted a specific exemption under the provisions of Section 18 of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, to the Arkansas State Plant Board for the use of Transform WG for control of tarnished plant bug in cotton. **Applicators must be in possession of the Transform WG Exemption Labeling at the time of application.** All applicable directions, restrictions, and precautions on the label for **Transform WG** and this notice must be followed.

Transform WG may be applied by ground or air at a rate of 1.5 to 2.75 oz. (0.047-0.086 lbs) of product per acre.

A maximum of 2 applications per acre per season.

A 14-day pre-harvest interval must be observed.

A maximum of 502,500 acres of cotton may be treated in the following counties: Ashley, Chicot, Clay, Craighead, Crittenden, Desha, Drew, Greene, Jackson, Jefferson, Lafayette, Lee, Lincoln, Lonoke, Mississippi, Monroe, Phillips, Poinsett, Prairie, Pulaski, and St. Francis.

Transform WG is highly toxic to bees. Do not apply this product while bees are actively visiting the treated area.

A report of each application made under the Specific Exemption must be filed with the Plant Board within 10 days. The following format is suggested for use by the applicator so the information required by the EPA will be provided: **THIS INFORMATION IS NOT TO BE REPORTED ON PINK FORM NUMBER 1007.**

REPORT OF TRANSFORM WG APPLICATION TO COTTON:

Applicator's Name:
Applicator's Address:
Farmer's Name:
Farmer's Address:
Acres Treated:
Pounds of Active Ingredient Applied per Acre:
Location of Field:
Time and Date of Application:

Reports may be mailed to: Arkansas State Plant Board, PO Box 1069, Little Rock, AR 72202-1069 or faxed to: (501)-312-7053.

Any adverse effects resulting from the use of Transform WG under this emergency exemption must be immediately reported to the Arkansas State Plant Board.

THIS EXEMPTION EXPIRES SEPTEMBER 30, 2012

Dow AgroSciences

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Transform™ WG

EPA Reg. No. 62719-XXX

Section 18 Specific Exemption

Control of Tarnished Plant Bug (*Lygus lineolaris*) in Cotton

(For Distribution and Use Only in the Counties of Ashley, Chicot, Clay, Craighead, Crittenden, Desha, Drew, Greene, Jackson, Jefferson, Lafayette, Lee, Lincoln, Lonoke, Mississippi, Monroe, Phillips, Poinsett, Prairie, Pulaski, and St. Francis in the State of Arkansas)

- **This Specific Exemption is effective June 1, 2012 through September 30, 2012.**
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Transform™ WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Transform WG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Transform WG.

Directions for Use

Pests and Application Rates:

Pests	Transform WG (oz/acre)
tarnished plant bug western tarnished plant bug	1.5 – 2.75 (0.047 – 0.086 lb ai/acre)

Application Timing: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Use a higher rate in the rate range for heavy pest populations. Two applications may be required for optimum tarnished plant bug control under high pest pressure or heavy immigration of plant bugs from other crops.

Restrictions:
- **Preharvest Interval:** Do not apply within 14 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 5 days apart.
- Do not make more than two applications per acre per year.
- Do not make more than one application in pre-bloom cotton.
- Do not apply more than a total of 5.5 oz of Transform WG (0.172 lb ai of sulfoxaflor) per acre per year.

™Trademark of Dow AgroSciences LLC

R396-001
Approved: __/__/__
File Symbol: 12AR02
Initial printing

(Base label):

Transform™ WG

Insecticide

For control of tarnished plant bug (*Lygus lineolaris*) in cotton

ATTENTION
- **Section 18 Specific Exemption**
- **For Distribution and Use Only in States with a Valid Section 18 Label**

This product cannot be used without a valid state-specific Section 18 label which must be in the possession of the user at the time of application.

Active Ingredient:
sulfoxaflor 50%
Other Ingredients 50%
Total 100%

Contains 50% active ingredient on a weight basis.

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Precautionary Statements

Hazard to Humans and Domestic Animals

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed

Do not get in eyes or on clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:
- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area.

There may be risk to honey bees through exposure to residues in pollen and nectar from flowers and from extra floral nectaries on cotton.

Advisory language: In order to minimize impact to honey bees when treating cotton, consider making applications in the early morning or early evening when bees are less likely to be visiting the treatment area.

To protect bees and other pollinators outside of the treatment area, this product should not be applied under conditions that could result in excessive drift to neighboring areas where bees are expected to be actively foraging.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Store in original container only.
Pesticide Disposal: Return any unopened, unused container of this unregistered product to the manufacturer or distributor, or dispose of it in accordance with Resource Conservation and Recovery Act regulations, following the expiration of the emergency exemption.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available

or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-XXX EPA Est. _____

™Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Weight ____

(Cover, shipping container):

Transform™ WG

Insecticide

For control of tarnished plant bug (*Lygus lineolaris*) in cotton

ATTENTION
- **Section 18 Specific Exemption**
- **For Distribution and Use Only in States with a Valid Section 18 Label**

This product cannot be used without a valid state-specific Section 18 label which must be in the possession of the user at the time of application.

Active Ingredient:
sulfoxaflor 50%
Other Ingredients 50%
Total 100%

Contains 50% active ingredient on a weight basis.

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-XXX EPA Est. _____

™Trademark of Dow AgroSciences LLC
Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Weight ____

(Page 1 through end):

Precautionary Statements

Hazard to Humans and Domestic Animals

DANGER

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed

Do not get in eyes or on clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area.

There may be risk to honey bees through exposure to residues in pollen and nectar from flowers and from extra floral nectaries on cotton.

Advisory language: In order to minimize impact to honey bees when treating cotton, consider making applications in the early morning or early evening when bees are less likely to be visiting the treatment area.

To protect bees and other pollinators outside of the treatment area, this product should not be applied under conditions that could result in excessive drift to neighboring areas where bees are expected to be actively foraging.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.
Pesticide Storage: Store in original container only.
Pesticide Disposal: Return any unopened, unused container of this unregistered product to the manufacturer or distributor, or dispose of it in accordance with Resource Conservation and Recovery Act regulations, following the expiration of the emergency exemption.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

Carefully read, understand and follow label use rates and restrictions. Apply the amount specified in the following tables with properly calibrated aerial or ground spray equipment. Prepare only the amount of

spray solution required to treat the measured acreage. The low rates may be used for light infestations of the target pests and the higher rates for moderate to heavy infestations. Transform™ WG insecticide may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of Transform WG per acre regardless of the spray volume used.

Use Precautions

Integrated Pest Management (IPM) Programs
Transform WG is recommended for IPM programs in labeled crops. Apply Transform WG when field scouting indicates target pest densities have reached the economic threshold, i.e., the point at which the insect population must be reduced to avoid economic losses beyond the cost of control. Other than reducing the target pest species as a food source, Transform WG does not have a significant impact on most parasitic insects or the natural predaceous arthropod complex in treated crops, including big-eyed bugs, ladybird beetles, flower bugs, lacewings, minute pirate bugs, damsel bugs, assassin bugs, predatory mites or spiders. The feeding activities of these beneficials will aid in natural control of other insects and reduce the likelihood of secondary pest outbreaks. If Transform WG is tank mixed with any insecticide that reduces its selectivity in preserving beneficial predatory insects, the full benefit of Transform WG in an IPM program may be reduced.

Insecticide Resistance Management (IRM)
Transform WG contains a Group 4C insecticide. Insect/mite biotypes with acquired resistance to Group 4C insecticides may eventually dominate the insect/mite population if Group 4C insecticides are used repeatedly in the same field or area, or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Transform WG or other Group 4C insecticides.

To delay development of insecticide resistance, the following practices are recommended:
- Avoid consecutive use of insecticides on succeeding generations with the same mode of action (same insecticide subgroup, 4C) on the same insect species.
- Consider tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use upon comprehensive IPM programs.
- Monitor treated insect populations in the field for loss of effectiveness.
- Do not treat seedling plants grown for transplant in greenhouses, shade houses, or field plots.
- Contact your local extension specialist, certified crop advisor, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Dow AgroSciences by calling 800-258-3033.

Mixing Directions

Application Rate Reference Table

Application Rate of Transform WG (oz/acre)	Active Ingredient Equivalent (lb ai/acre)
1.5	0.047
1.75	0.055
2.75	0.086

Transform WG – Alone
Fill the spray tank with water to about 1/2 of the required spray volume. Start agitation and add the required amount of Transform WG. Continue agitation while mixing and filling the spray tank to the required spray volume. Maintain sufficient agitation during application to ensure uniformity of the spray mix. Do not allow water or spray mixture to back-siphon into the water source.

Transform WG - Tank Mix
When tank mixing Transform WG with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. If foliar fertilizers are used, the jar test should be repeated with each batch of fertilizer utilizing the mixing water source. Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes: Fill the spray tank with water to 1/4 to 1/3 of the required spray volume. Start agitation. Add different formulation types in the order indicated below, allowing time for complete dispersion and mixing after addition of each product. Allow extra dispersion and mixing time for dry flowable products.

Add different formulation types in the following order:
1. Transform WG and other water dispersible granules
2. Wettable powders
3. Suspension concentrates and other liquids

Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add:
4. Emulsifiable concentrates and water-based solutions
5. Spray adjuvants, surfactants and oils
6. Foliar fertilizers

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose.

Premixing: Dry and flowable formulations may be premixed with water (slurried) and added to the spray tank through a 20 to 35 mesh screen. This procedure assures good initial dispersion of these formulation types.

Application Directions

Proper application techniques help ensure thorough spray coverage and correct dosage for optimum insect control. Apply Transform WG as a foliar spray at the rate indicated for target pest. The following directions are provided for ground and aerial application of Transform WG. Attention should be given to sprayer speed and calibration, wind speed, and foliar canopy to ensure adequate spray coverage.

Spray Drift Management
Wind: To reduce off-target drive and achieve maximum performance, apply when wind velocity favors on-target product deposition.

Temperature Inversions: Do not make ground or aerial applications during a temperature inversion. Temperature inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size: Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASABE (S-572.1) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size except where indicated for specific crops.

Ground Application

To avoid drift for groundboom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy. Shut off the sprayer when turning at row ends. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind directions are toward the aquatic area.

Aerial Application
Apply in a minimum spray volume of 3 gallons per acre. Mount the spray boom on the aircraft so as to minimize drift caused by wing tip or rotor vortices. Use the minimum practical boom length and do not exceed 75% of the wing span or 80% of the rotor diameter. Flight speed and nozzle orientation must be considered in determining droplet size. Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Spray Adjuvants
The addition of agricultural adjuvants to sprays of Transform WG may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Transform WG at specified rates for a registered use.

Crop	Re-Planting Interval
cotton	no restrictions
all other food and feed crops	360 days

Application Rates

Refer to state-specific Section 18 label for approved uses, application timing and restrictions.

This product cannot be used without a valid state-specific Section 18 label. The label must be in the possession of the user at the time of application. The Section 18 label can be obtained from state and/or county authorities where it is valid. Please contact your Dow AgroSciences representative for further information.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®™Trademark of Dow AgroSciences LLC
Accepted 06/01/12